

1/2 sister to Ferguson  
Kate Burke & Clark  
Acker. Murdoch

Mervyn Mowbray  
Vilate 11 111111

Ed McMillan  
Pop Raymond Mgr

= star of dawn  
Sarah Crook

F = John "Jack"  
Carlile Smith

Emma Amelia  
Deber Merc

Song of  
Africa 1912

part of 1912

the School

Deber Merc

## UTAH INFLUENZA IMMUNIZATION PROJECT

### INFLUENZA VACCINE REACTION INFORMATION

You have just received influenza vaccine. A small percentage of persons receiving this vaccine may experience one or more of the following symptoms: redness and tenderness at the injection site, fever, (usually 101° or less), chills, nausea, loss of appetite, muscle ache, joint pain, headache, or fatigue. These reactions are usually short lived, lasting less than 48 hours, but local reactions at the place of injection may persist for several days. If you experience severe or prolonged reaction, consult your physician or call your local health department at



## UTAH INFLUENZA IMMUNIZATION PROJECT

### INFLUENZA VACCINE REACTION INFORMATION

You have just received influenza vaccine. A small percentage of persons receiving this vaccine may experience one or more of the following symptoms: redness and tenderness at the injection site, fever, (usually 101° or less), chills, nausea, loss of appetite, muscle ache, joint pain, headache, or fatigue. These reactions are usually short lived, lasting less than 48 hours, but local reactions at the place of injection may persist for several days. If you experience severe or prolonged reaction, consult your physician or call your local health department at

Ray National  
Booth Report

How Much  
1896 27 yr

9 yrs

~~1905~~

The Washington

old How  
1903

## UTAH INFLUENZA IMMUNIZATION PROJECT

### INFLUENZA VACCINE REACTION INFORMATION

You have just received influenza vaccine. A small percentage of persons receiving this vaccine may experience one or more of the following symptoms: redness and tenderness at the injection site, fever, (usually 101° or less), chills, nausea, loss of appetite, muscle ache, joint pain, headache, or fatigue. These reactions are usually short lived, lasting less than 48 hours, but local reactions at the place of injection may persist for several days. If you experience severe or prolonged reaction, consult your physician or call your local health department at